UBE2L3

Cat. No.	SSB-CE0020
Lot. No.	163060020

UBE2L3

UBE2L3 is an E2 ubiquitin conjugating enzyme and accepts ubiquitin from an E1 activating enzyme via an active site cysteine. The mechanism of ubiquitin transfer involves the breaking of a E1-Ub thioester linkage, followed by a reformation of a UBE2L3-Ub thioester. UBE_{2L3} transfers active ubiquitin molecules via a final transthiolation reaction to a cysteine residue of HECT class or RBR class E3 ligases. UBE2L3 plays a critical role in the activation of N-kb signaling by working in conjugation with the RBR E3 ligase, LUBAC (linear ubiquitin assembly complex), to ubiquitinate NEMO of the IkB complex, which than facilitates the downstream immune response. This recombinant UBE₂L₃ is expressed in E.coli with an N-terminal polyhistidine tag.



South Bay Bío

Product Information

Quantity: 100µg Molecular Weight: 18 kDa

Concentration: 50 µM, 0.9mg/mL

Purity: >95% by SDS-PAGE

Storage Buffer: HEPES pH 7.5, 150mM NaCl, 10% glycerol, 2mM TCEP

Storage: -80C, Avoid multiple freeze / thaw

Quality Control and Performance Data



UBE2L3 SDS-PAGE. From left to right, increasing amounts of UBE2L3 loaded onto a 4-20% SDS-PAGE gel, stained with coomassie brillant blue. Purity is > 95%.

For Research Use Only, Not For Use In Humans.

www.southbaybio.com

Contact: info@southbaybio.com 5941 Optical Ct, Suite 229 San Jose, CA 95138 USA

UBE2L3



Thioester Activity Assay. UBE2L3 forms a thioester with UB in an ATP dependent manner, and the bond can be reduced with addition of excess DTT. The UBE2L3 is active.

For Research Use Only, Not For Use In Humans.

www.southbaybio.com

Contact: info@southbaybio.com 5941 Optical Ct, Suite 229 San Jose, CA 95138 USA