

K33-linked Di-Ubiquitin

Cat. No. SBB-UP0066
Lot. No. 181940066



South Bay Bio

K33 Di-Ubiquitin

The array of cellular processes initiated and regulated by ubiquitin has been partially explained by the structural diversity of differently linked ubiquitin chains. In a ubiquitin chain, ubiquitin moieties can be conjugated through one of their lysine residues (K6, K11, K27, K29, K33, K48 and K63) or the N-terminal methionine residue (M1), offering countless possibilities to assemble a specific polymer. Ubiquitin molecules can also be modified by other post-translational modifications, including acetylation and phosphorylation, adding another layer of ubiquitin signal regulation and diversification.

K33-linked ubiquitin chains are associated with negative regulation of both T-cell antigen receptor (TCR) and AMP-activated protein kinase (AMPK)-related protein kinases. K33 linkages have also been implicated in post-Golgi protein trafficking. This K33 linked di-ubiquitin was enzymatically conjugated, and purified via liquid chromatography.



References

- 1) Dikic, I., Wakatsuki, S., & Walters, K. J. (2009). Ubiquitin-binding domains – from structures to functions. *Nature Reviews Molecular Cell Biology*, 10(10), 659-671. <https://doi.org/10.1038/nrm2767>
- 2) Akutsu, M., Dikic, I., & Bremm, A. (2016). Ubiquitin chain diversity at a glance. *Journal of Cell Science*, 129(5), 875-880. <https://doi.org/10.1242/jcs.183954>

Product Information

Quantity: 25 µg **Molecular Weight:** 17 kDa


Concentration: 58 µM, 1 mg/mL

Purity: >95% by SDS-PAGE

Storage Buffer: 50 mM HEPES pH 7.5

Storage: -80C, Avoid multiple freeze / thaw

Quality Control and Performance Data



K33-Linked Di-Ubiquitin SDS-PAGE. From left to right, increasing amounts of di-ubiquitin were loaded onto a 10-20% SDS-PAGE gel, stained with Coomassie brilliant blue. Purity is >95%.

For Research Use Only, Not For Use In Humans.

www.southbaybio.com

Contact:
info@southbaybio.com

5941 Optical Ct, Suite 229
San Jose, CA 95138 USA