## K48 Tri-Ubiquitin

Cat. No. SBB-UP0171 Lot. No. 243020171



The array of cellular processes initiated and regulated by ubiquitin has been partially explained by the structural diversity of differently linked ubiquitin chains. In a ubiquitin chain, ubiquitin moieties can be conjugated through one of their lysine residues (K6, K11, K27, K29, K33, K48 and K63) or the N-terminal methionine residue (M1), offering countless possibilities to assemble a specific polymer. Ubiquitin molecules can also be modified by other post-translational modifications, including acetylation and phosphorylation, adding another layer of ubiquitin signal regulation and diversification.

K48-polyubiquitin chains are the most abundant linkage in cells and thought to be the major signal for proteasome-mediated degradation. Quantitative MS analyses of intracellular ubiquitin linkages reveled K48-polyubiquitin linkages rapidly accumulate when cells are treated with the proteasome inhibitor MG132. This K48-linked Triubiquitin is enzymatically conjugated, and purified via liquid chromatography.

## References

- 1) Dikic, I., Wakatsuki, S., & Walters, K. J. (2009). Ubiquitin-binding domains from structures to functions. Nature Reviews Molecular Cell Biology, 10(10), 659–671. https://doi.org/10.1038/nrm2767
- 2) Akutsu, M., Dikic, I., & Bremm, A. (2016). Ubiquitin chain diversity at a glance. Journal of Cell Science, 129(5), 875–880. https://doi.org/10.1242/jcs.183954



## **Product Information**

Quantity: 25 μg Molecular Weight: 26 kDa


Concentration: 39 µM, 1 mg/mL

Purity: >98% by SDS-PAGE

Storage Buffer: 1x PBS

Storage: -80C, Avoid multiple freeze / thaw

## **Quality Control and Performance Data**



K48-Linked Tri-Ubiquitin SDS-PAGE. From left to right, increasing amounts of tri-ubiquitin were loaded onto a 10-20% SDS-PAGE gel, stained with Coomassie brillant blue. Purity is > 98%.

For Research Use Only, Not For Use In Humans.

www.south-bay-bio.com