





**PRODUCT DATA SHEET** )

Page 1/3

## MPLA from S. minnesota R595 (Re) TLRpure<sup>™</sup> Sterile Solution

Source	Monophosphoryl Lipid A [MPLA] derived from S. minnesota R595 (Re) TLRpure™ LPS
Concentration	Img/ml (0.5mg/ml for 250µg size) stabilised in sterile, double-distilled water (ddWater) without any additives
TLRpure™	No detectable TLR4 independent activity: standardised potent TLR4-specific agonist
Purity	Ultrapure. No detectable DNA, RNA and protein traces.
Purification Method	R-type (mutant/rough) LPS was isolated by a phenol-chloroform-petroleum-ether method. Semi-purified LPS was subjected to further re-extraction cycles and ultracentrifugation steps, extensively electrodialysed before converted to its uniform sodium salt form to yield TLRpure <sup>™</sup> LPS, from which MPLA was generated by acid hydrolysis.
Sterility	Filter method: according to Ph. Eur. 9. Passed according to specification: • No growth in Thioglycolate medium at 30-35°C after 14 days. • No growth in Soybean Casein Digest Broth (TSB) at 20-25°C after 14 days.
Endotoxin Content	Bacterial Endotoxin Test (kinetic turbidimetric LAL method) according to Ph. Eur. 9. Endotoxin Content: >1,000,000 [EU/ml].
Appearance	Colourless, clear to opaque aqueous solution
Handling	Keep sterile. Prepare working dilutions from pre-warmed (~40°C) MPLA stock solution just prior to use. Ready-made solution is cell culture-grade. Do not pre-dilute in buffer (e.g. PBS) as this will lead to precipitation of MPLA. To yield a 100µg/ml (1,000-100x) stock solution add 100µl of MPLA to 900µl endotoxin-free and sterile ddWater (Cat. No.: IAX-900-002) (not PBS) and mix well.
Activity	Optimal concentration is dependent upon cell type, species, desired activation and analysis: 0.1-1.0µg/ml <i>in vitro</i> and 5-15mg/kg <i>in vivo</i> in animal rodent models. Does not activate any TLR other than TLR4 as tested up to 1µg/ml in relevant cellular systems (mouse macrophages).
Shipping	Ambient
Storage	2-8°C
Stability	2 years after receipt (unopened and as supplied). Diluted solutions are stable for 12 hours at 2-8°C.
MSDS	Available on request

Document No.: IAX-100-002 | Version: 1.9 | Issue Date: 12/02/2023

DISCLAIMER: THIS PRODUCT IS NOT INTENDED OR APPROVED FOR HUMAN, DIAGNOSTICS OR VETERINARY USE. USE OF THIS PRODUCT FOR HUMAN OR ANIMAL TESTING MAY BE EXTREMELY HAZARDOUS AND MAY RESULT IN DISEASE, SEVERE INJURY, OR DEATH. THIS PRODUCT IS FOR RESEARCH USE ONLY (RUO). MATERIAL SAFETY DATA: This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, inhale or get into the blood stream. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Access to this material must be restricted to personnel, who is appropriately experienced, qualified, competent and properly trained to use it. Material Safety Data Sheet is available upon request.



Innaxon United Kingdom Email: info@innaxon.com Website: www.innaxon.co.uk



## PRODUCT DATA SHEET

**Page** 2/3

Cat. No.: IAX-100-002	Lot. No.:
Product Information	<ul> <li>MPLA has been generated by acid hydrolysis from TLRpure<sup>™</sup> LPS purified according to an optimised and proprietary extraction and purification protocol, but based upon the methods published by Galanos, et al. (laboratory of Westphal and Lüderitz, Freiburg, Germany).</li> <li>TLRpure<sup>™</sup> LPS lacks any detectable bacterial, (lipo-)protein, RNA or DNA or other TLR-stimulating activity due to its ultra-purified formulation. Its unique potency and purity are quality controlled using a physiological system of primary innate immune cells and a relevant biological cytokine expression read-out.</li> <li>All immunological activity of the MPLA is exclusively dependent upon the presence of TLR4 as determined by the use of the corresponding control cells, where TLR4 has been genetically deleted or missing (from TLR4 deficient also called TLR4 knock-out KO mice).</li> <li>TLRpure<sup>™</sup> MPLA convenient ready-made stabilised solution makes it the reagent of choice for in vitro as well as in vivo experiments for superior reproducible and comparable results.</li> <li>Compared to MPLA derived from conventional (semi-purified) LPS preparations, this product is derived from low yield TLRpure<sup>™</sup> LPS produced on an industrial fermentation scale under precisely controlled growth conditions to yield large batch sizes, allowing custom formulations/packaging.</li> </ul>
Product Specific Reference	es [1] A new method for the extraction of R lipopolysaccharides. Galanos C, et al. Eur. J. Biochem. (1969); 9:245
	[2] Endotoxic properties of chemically synthesized lipid A part structures. Comparison of synthetic lipid A precursor and synthetic analogues with biosynthetic lipid A precursor and free lipid A. Galanos C, et al. Eur. J. Biochem. (1984); 140:221
	[3] The Biosynthetic Monophosphoryl Lipid A Enhances the Therapeutic Outcome of Antibiotic Therapy in Pneumococcal Pneumonia. Casilag F, et al. ACS Infect. Dis. (2021); 7:2164-2175
	[4] Synthetic Glycolipids as Molecular Vaccine Adjuvants: Mechanism of Action in Human Cells and In Vivo Activity. Facchini FA, et al. J. Med. Chem. (2021); 64:12261
	[5] Synthetic glycolipid-based TLR4 antagonists negatively regulate TRIF-dependent TLR4 signalling in human macrophages. Palmer C, Facchini F A, Jones R PO, Neumann F, Peri F, Pirianov G. Innate Immunity (2021); 27:275-284
	[6] Overcoming Challenges in Chemical Glycosylation to Achieve Innovative Vaccine Adjuvants Possessing

Document No.: IAX-100-002 | Version: 1.9 | Issue Date: 12/02/2023

DISCLAIMER: THIS PRODUCT IS NOT INTENDED OR APPROVED FOR HUMAN, DIAGNOSTICS OR VETERINARY USE. USE OF THIS PRODUCT FOR HUMAN OR ANIMAL TESTING MAY BE EXTREMELY HAZARDOUS AND MAY RESULT IN DISEASE, SEVERE INJURY, OR DEATH. THIS PRODUCT IS FOR RESEARCH USE ONLY (RUO). MATERIAL SAFETY DATA: This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, inhale or get into the blood stream. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Access to this material must be restricted to personnel, who is appropriately experienced, qualified, competent and properly trained to use it. Material Safety Data Sheet is available upon request.



Innaxon United Kingdom Email: info@innaxon.com Website: www.innaxon.co.uk



## PRODUCT DATA SHEET

**Page** 3/3

<ul> <li>Activation of cells by LPS is mediated by the Toll-like receptor 4 (TLR4), a member of the highly conserved protein family of TLRs, which are specialised in the recognition of microbial components. In mice, defects in TLR4 result in LPS unresponsiveness.</li> <li>For optimal interaction with LPS, TLR4 requires association with myeloid differentiation protein (LBP). This mechanism is believed to be generally true for LPS signaling, Re-form LPS and lipid A, but not S-form LPS, are capable of inducing TNF-a responses also in the absence of CD14.</li> <li>LPS, synthesized by most wild-type (WT) Gram-negative bacteria (S-form LPS), consists of three regions, the O-polysaccharide and the lipid A, which harbors the endotoxic activity of the entire molecule. R-form LPS synthesized by the so-called rough (R) mutants of Gram-negative bacteria lacks the O-specific chain. Furthermore, the core-oligosaccharide mutan tbelongs, Monophosphoryl Lipid A (MPLA) represents a detoxified derivative of Lipid A and constitutes an important adjuvant in prophylactic and therapeutic vaccines.</li> <li>References</li> <li>[1] <i>R-form LPS, the master key to the activation of TLR4/MD-2-positive cells</i>. Huber M, et al. Eur. J. Immunol. (2006); 36:701</li> <li>[2] CD14 is required for MyD88-independent LPS signaling, Jang Z, Georgel P, Du X, Shamel L, Sovarta S, Mudd S, Huber M, Kalis C, Keck S, Galanos C, Freudenberg M, Beutler B. Nat. Immunol. (2005); 6:565</li> <li>[3] Defective LPS signaling in G3HIHelg and G578L/105CCr mice mutations in TIr4 gene. Poltorak A, He X, Smirmova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciard-Castagnoli P, Layton B, Beutler B. Science (1998); 28:22085</li> <li>[4] Structural relationship of Salmonella 0 and R antigens. Lüderitz O, Galanos C, et al. Ann. N.Y. Acad. Sci. (1966); 133:349</li> <li>[5] Preparation and properties of antisera against the lipid A component of bacterial lipopolysaccharides. Galanos C, et al.</li></ul>	Cat. No.: IAX-100-002	Lot. No.:
<ul> <li>[1] R-form LPS, the master key to the activation of TLR4/MD-2-positive cells. Huber M, et al. Eur. J. Immunol. (2006); 36:701</li> <li>[2] CD14 is required for MyD88-independent LPS signaling. Jiang Z, Georgel P, Du X, Shamel L, Sovath S, Mudd S, Huber M, Kalis C, Keck S, Galanos C, Freudenberg M, Beutler B. Nat. Immunol. (2005); 6:565</li> <li>[3] Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. Science (1998); 282:2085</li> <li>[4] Structural relationship of Salmonella 0 and R antigens. Lüderitz O, Galanos C, et al. Ann. N.Y. Acad. Sci. (1966); 133:349</li> <li>[5] Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides. Galanos C, et al. Lur. J. Biochem. (1971); 24:116</li> <li>[6] Lipid A: chemical structure and biological activity. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17</li> <li>[7] Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of</li> </ul>	General Information	<ul> <li>highly conserved protein family of TLRs, which are specialised in the recognition of microbial components. In mice, defects in TLR4 result in LPS unresponsiveness.</li> <li>For optimal interaction with LPS, TLR4 requires association with myeloid differentiation protein 2 (MD-2). According to current consensus activation of TLR4 is preceded by the transfer of LPS to membrane-bound (m) or soluble (s) CD14 by LPS-binding protein (LBP). This mechanism is believed to be generally true for LPS signaling. Re-form LPS and lipid A, but not S-form LPS, are capable of inducing TNF-α responses also in the absence of CD14.</li> <li>LPS, synthesized by most wild-type (WT) Gram-negative bacteria (S-form LPS), consists of three regions, the O-polysaccharide chain, which is made up of repeating oligosaccharide units, the core oligosaccharide and the lipid A, which harbors the endotoxic activity of the entire molecule. R-form LPS synthesized by the so-called rough (R) mutants of Gram-negative bacteria lacks the O-specific chain. Furthermore, the core-oligosaccharide may be present in different degrees of completion, depending on the class (Ra to Re) to which the mutant belongs Monophosphoryl Lipid A (MPLA) represents a detoxified derivative of Lipid A and constitutes</li> </ul>
<ul> <li>Eur. J. Immunol. (2006); 36:701</li> <li>[2] <i>CD14 is required for MyD88-independent LPS signaling</i>. Jiang Z, Georgel P, Du X, Shamel L, Sovath S, Mudd S, Huber M, Kalis C, Keck S, Galanos C, Freudenberg M, Beutler B. Nat. Immunol. (2005); 6:565</li> <li>[3] <i>Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene</i>. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. Science (1998); 282:2085</li> <li>[4] <i>Structural relationship of Salmonella 0 and R antigens</i>. Lüderitz O, Galanos C, et al. Ann. N.Y. Acad. Sci. (1966); 133:349</li> <li>[5] <i>Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides</i>. Galanos C, et al. Eur. J. Biochem. (1971); 24:116</li> <li>[6] <i>Lipid A: chemical structure and biological activity</i>. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17</li> <li>[7] <i>Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of</i></li> </ul>	References	
<ul> <li>Sovath S, Mudd S, Huber M, Kalis C, Keck S, Galanos C, Freudenberg M, Beutler B. Nat. Immunol. (2005); 6:565</li> <li>[3] Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. Science (1998); 282:2085</li> <li>[4] Structural relationship of Salmonella 0 and R antigens. Lüderitz O, Galanos C, et al. Ann. N.Y. Acad. Sci. (1966); 133:349</li> <li>[5] Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides. Galanos C, et al. Eur. J. Biochem. (1971); 24:116</li> <li>[6] Lipid A: chemical structure and biological activity. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17</li> <li>[7] Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of</li> </ul>		
<ul> <li>He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. Science (1998); 282:2085</li> <li>[4] Structural relationship of Salmonella 0 and R antigens. Lüderitz O, Galanos C, et al. Ann. N.Y. Acad. Sci. (1966); 133:349</li> <li>[5] Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides. Galanos C, et al. Eur. J. Biochem. (1971); 24:116</li> <li>[6] Lipid A: chemical structure and biological activity. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17</li> <li>[7] Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of</li> </ul>		Sovath S, Mudd S, Huber M, Kalis C, Keck S, Galanos C, Freudenberg M, Beutler B.
<ul> <li>Acad. Sci. (1966); 133:349</li> <li>[5] Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides. Galanos C, et al. Eur. J. Biochem. (1971); 24:116</li> <li>[6] Lipid A: chemical structure and biological activity. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17</li> <li>[7] Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of</li> </ul>		He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C,
Galanos C, et al. Eur. J. Biochem. (1971); 24:116 [6] Lipid A: chemical structure and biological activity. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17 [7] Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of		
[7] Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of		
		[6] Lipid A: chemical structure and biological activity. Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17

**Document No.:** IAX-100-002 | **Version:** 1.9 | **Issue Date:** 12/02/2023

DISCLAIMER: THIS PRODUCT IS NOT INTENDED OR APPROVED FOR HUMAN, DIAGNOSTICS OR VETERINARY USE. USE OF THIS PRODUCT FOR HUMAN OR ANIMAL TESTING MAY BE EXTREMELY HAZARDOUS AND MAY RESULT IN DISEASE, SEVERE INJURY, OR DEATH. THIS PRODUCT IS FOR RESEARCH USE ONLY (RUO). MATERIAL SAFETY DATA: This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, inhale or get into the blood stream. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Access to this material must be restricted to personnel, who is appropriately experienced, qualified, competent and properly trained to use it. Material Safety Data Sheet is available upon request.